arXiv:1010.5936v1 [math.DG] 28 Oct 2010

Spinor-generators of compact exceptional Lie groups F_4, E_6 and E_7

Takashi Miyasaka, Osamu Shukuzawa and Ichiro Yokota

1 Introduction

We know that any element A of the group $SO(3)$ can be represented as

$$A = A_1A_2{A_1}', \quad A_1, {A_1}' \in SO_1(2), A_2 \in SO_2(2)$$

where $SO_k(2) = \{A \in SO(3) \,|\, A\boldsymbol{e}_k = \boldsymbol{e}_k\}(k = 1, 2)$ ([1]). In the present paper, we shall show firstly that the similar results hold for the groups $SU(3)$, and $Sp(3)$ (Theorem 1). Secondly, we shall show that any element α of the simply connected compact Lie group F_4 (resp. E_6) can be represented as

$$\alpha = \alpha_1\alpha_2{\alpha_1}', \qquad \begin{array}{c} \alpha_1, {\alpha_1}' \in Spin_1(9), \alpha_2 \in Spin_2(9) \\ (\text{resp.}\alpha_1, {\alpha_1}' \in Spin_1(10), \alpha_2 \in Spin_2(10)) \end{array}$$

where $Spin_k(9) = \{\alpha \in F_4 \,|\, \alpha E_k = E_k\}$ (resp. $Spin_k(10) = \{\alpha \in E_6 \,|\, \alpha E_k = E_k\}$ (Theorem 5 (resp. Theorem 7))). Lastly, we shall show that any element α of the simply connected compact Lie group E_7 can be represented as

$$\alpha = \alpha_1\alpha_2{\alpha_1}'{\alpha_2}'{\alpha_1}'', \quad \alpha_1, {\alpha_1}', {\alpha_1}'' \in Spin_1(12), \alpha_2, {\alpha_2}' \in Spin_2(12)$$

where $Spin_k(12) = \{\alpha \in E_7 \,|\, \alpha\kappa_k = \kappa_k\alpha, \alpha\mu_k = \mu_k\alpha\}$ (Theorem 10).

In this paper we follow the notation of [2].

2 Spinor-generators of the groups $SO(3), SU(3)$ and $Sp(3)$

Let $\boldsymbol{H}$ be the quaternion field with basis 1, $\boldsymbol{i}, \boldsymbol{j}$ and $\boldsymbol{k}$ over $\boldsymbol{R}$. Then we can express each element $a = a_0 + a_1\boldsymbol{i} + a_2\boldsymbol{j} + a_3\boldsymbol{k} \in \boldsymbol{H}$ in the following polar form

$$a = r(\cos\theta + u\sin\theta), \quad u^2 = -1\,(u \in \boldsymbol{H}),\ r = |a| = \sqrt{\sum_{k=0}^{3} {a_k}^2}, \theta \in \boldsymbol{R}.$$

Hereafter, we briefly denote by $re^{u\theta}$ an element $r(\cos\theta + u\sin\theta)$ after the model of complex numbers.

The classical groups $SO(n), SU(n)$ and $Sp(n)$ are respectively defined by

$$\begin{aligned}
&SO(n) = \{A \in M(n, \boldsymbol{R}) \mid {}^tAA = E, \det A = 1\},\\
&SU(n) = \{A \in M(n, \boldsymbol{C}) \mid A^*A = E, \det A = 1\},\\
&Sp(n) = \{A \in M(n, \boldsymbol{H}) \mid A^*A = E\}
\end{aligned}$$

where we follow the usual convention for matrices : $M(n, K)$ (= the set of square matrices of order n with coefficients in $K = \boldsymbol{R}, \mathbf{C}$ or $\boldsymbol{H}$), ${}^tA, A^*(= \overline{{}^tA}), E$ (= the unit matrix) and det (= the determinant).

Theorem 1. (1) *Any element* $A \in SO(3)$ *can be represented as*

$$A = A_1A_2{A_1}', \quad A_1, {A_1}' \in SO_1(2), A_2 \in SO_2(2)$$

where $SO_k(2) = \{A \in SO(3) \mid A\boldsymbol{e}_k = \boldsymbol{e}_k\} \cong Spin(2)\,(k = 1, 2), \boldsymbol{e}_1 = {}^t(1, 0, 0), \boldsymbol{e}_2 = {}^t(0, 1, 0)$.

(2) *Any element* $A \in SU(3)$ *can be represented as*

$$A = A_1A_2{A_1}', \quad A_1, {A_1}' \in SU_1(2), A_2 \in SU_2(2)$$

where $SU_k(2) = \{A \in SU(3) \mid A\boldsymbol{e}_k = \boldsymbol{e}_k\} \cong Spin(3)\,(k = 1, 2)$.

(3) *Any element* $A \in Sp(3)$ *can be represented as*

$$A = A_1A_2{A_1}', \quad A_1, {A_1}' \in Sp_1(2), A_2 \in Sp_2(2)$$

where $Sp_k(2) = \{A \in Sp(3) \mid A\boldsymbol{e}_k = \boldsymbol{e}_k\} \cong Spin(3)\,(k = 1, 2)$.

Proof It suffices to prove (3), because we can reduce (1) and (2) to the particular case of (3) in the proof below. First, for a given element $A \in Sp(3)$, suppose $A\boldsymbol{e}_1 = {}^t(a_1, a_2, a_3), a_2 \neq 0\,(a_k \in \boldsymbol{H}\,(k = 1, 2, 3))$. Then there exists an element $u \in \boldsymbol{H}$ satisfying $u^2 = -1$ and a real number $\alpha \in \boldsymbol{R}$ such that $a_3a_2^{-1} = (|a_3|/|a_2|)e^{u\alpha}$. Choose $\theta \in \boldsymbol{R}$ such that $\cot\theta = |a_3|/|a_2|$ and set

$$B_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{u\alpha/2}\cos\theta & -e^{-u\alpha/2}\sin\theta \\ 0 & e^{u\alpha/2}\sin\theta & e^{-u\alpha/2}\cos\theta \end{pmatrix} \in Sp_1(2).$$

Then we get

$$B_1A\boldsymbol{e}_1 = {}^t(b_1, 0, b_3), \quad b_1, b_3 \in \boldsymbol{H}.$$

Next suppose $b_3 \neq 0$. Then there exists an element $v \in \boldsymbol{H}$ satisfying $v^2 = -1$ and a real number $\beta \in \boldsymbol{R}$ such that $b_1{b_3}^{-1} = (|b_1|/|b_3|)e^{v\beta}$. Choose $\varphi \in \boldsymbol{R}$ such that $\cot\varphi = -|b_1|/|b_3|$ and set

$$B_2 = \begin{pmatrix} e^{-v\beta/2}\cos\varphi & 0 & -e^{v\beta/2}\sin\varphi \\ 0 & 1 & 0 \\ e^{-v\beta/2}\sin\varphi & 0 & e^{v\beta/2}\cos\varphi \end{pmatrix} \in Sp_2(2).$$

Then we get

$$B_2B_1A\boldsymbol{e}_1 = {}^t(c_1, 0, 0), \quad c_1 \in \boldsymbol{H}.$$

Since $|c_1| = 1$, we can say $c_1 = e^{w\gamma}$ ($w^2 = -1, w \in \boldsymbol{H}, \gamma \in \boldsymbol{R}$). Set

$$B_2{}' = \begin{pmatrix} e^{-w\gamma} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{w\gamma} \end{pmatrix} \in Sp_2(2).$$

Then, since it follows $B_2{}'B_2B_1A\boldsymbol{e}_1 = \boldsymbol{e}_1$, *i.e.*, $B_2{}'B_2B_1A \in Sp_1(2)$, we can set $B_2{}'B_2B_1A = B_1{}' \in Sp_1(2)$. This implies

$$A = A_1A_2A_1{}', \quad A_1, A_1{}' \in Sp_1(2),\ A_2 \in Sp_2(2).$$

3 Some elements of $Spin_k(9), Spin_k(10)$ and $Spin_k(12)$.

As for the definitions of $Spin_k(9), Spin_k(10)$ and $Spin_k(12) (k = 1, 2)$, see Sections 4, 5 and 6.

Lemma 2 (Section 4 and [2]). (1) *Let* $\alpha_1(a)$ *be the mapping* $\alpha(a)$ *defined in* [2] *Lemma* 2.(1). *Then* $\alpha_1(a)$ *belongs to* $Spin_1(9) \subset Spin_1(10) \subset Spin_1(12)$.

(2) *For* $a \in \mathfrak{C}, a \neq 0$, *let* $\alpha_2(a) : \mathfrak{J} \to \mathfrak{J}$ *be the mapping defined by changing all of the indices from* k *to* $k+1$ (*index modulo* 3) *in the definition of* $\alpha(a)$ *of* [2] *Lemma* 2.(1). *Then* $\alpha_2(a)$ *belongs to* $Spin_2(9) \subset Spin_2(10) \subset Spin_2(12)$.

Lemma 3 (Section 5 and [2]). (1) *Let* $\beta_1(a)$ *be the mapping* $\beta(a)$ *defined in* [2] *Lemma* 2.(2). *Then* $\beta_1(a)$ *belongs to* $Spin_1(10) \subset Spin_1(12)$.

(2) *For* $a \in \mathfrak{C}, a \neq 0$, *let* $\beta_2(a) : \mathfrak{J}^C \to \mathfrak{J}^C$ *be the mapping defined by changing all of the indices from* k *to* $k+1$ (*index modulo* 3) *in the definition of* $\beta(a)$ *of* [2] *Lemma* 2.(2). *Then* $\beta_2(a)$ *belongs to* $Spin_2(10) \subset Spin_2(12)$.

Lemma 4 (Section 6 and [2]). (1) *Let* $\gamma_1(a)$ *be the mapping* $\gamma(a)$ *defined in* [2] *Lemma* 3.(1). *Then* $\gamma_1(a)$ *belongs to* $Spin_1(12)$.

(2) *For* $a \in \mathfrak{C}, a \neq 0$, *let* $\gamma_2(a) : \mathfrak{P}^C \to \mathfrak{P}^C$ *be the mapping defined by changing all of the indices from* k *to* $k+1$ (*index modulo* 3) *in the definition of* $\gamma(a)$ *of* [2] *Lemma* 3.(1). *Then* $\gamma_2(a)$ *belongs to* $Spin_2(12)$.

(3) *Let* $\delta_1(a)$ *be the mapping* $\delta(a)$ *defined in* [2] *Lemma* 3.(2). *Then* $\delta_1(a)$ *belongs to* $Spin_1(12)$.

(4) *For* $a \in \mathfrak{C}, a \neq 0$, *let* $\delta_2(a) : \mathfrak{P}^C \to \mathfrak{P}^C$ *be the mapping defined by changing all of the indices from* k *to* $k+1$ (*index modulo* 3) *in the definition of* $\delta(a)$ *of* [2] *Lemma* 3.(2). *Then* $\delta_2(a)$ *belongs to* $Spin_2(12)$.

4 $Spin(9)$-generatos of the group F_4

The simply connected compact Lie group F_4 is given by

$$F_4 = \{\alpha \in \mathrm{Iso}_R(\mathfrak{J}) \,|\, \alpha(X \times Y) = \alpha X \times \alpha Y\}.$$

The group F_4 has subgroups

$$Spin_k(9) = \{\alpha \in F_4 \,|\, \alpha E_k = E_k\} \quad (k = 1, 2),$$

where $E_1 = (1, 0, 0; 0, 0, 0), E_2 = (0, 1, 0; 0, 0, 0) \in \mathfrak{J}$, which is isomorphic to the usual spinor group $Spin(9)$ ([2],[3]).

Theorem 5. *Any element $\alpha \in F_4$ can be represented as*

$$\alpha = \alpha_1\alpha_2\alpha_1{}', \quad \alpha_1, \alpha_1{}' \in Spin_1(9), \alpha_2 \in Spin_2(9).$$

Proof For a given element $\alpha \in F_4$, it suffices to show that there exist $\alpha_1 \in Spin_1(9)$ and $\alpha_2 \in Spin_2(9)$ such that $\alpha_2\alpha_1\alpha E_1 = E_1$. Now, for $\alpha E_1 = (\xi_1, \xi_2, \xi_3; x_1, x_2, x_3) = X_0$, choose $a \in \mathfrak{C}$ such that $(a, x_1) = 0, |a| = \pi/4$, and define $\alpha_1(a) \in Spin_1(9)$ of Lemma 2.(1). Then we get

$$\alpha_1(a)X_0 = (\xi_1{}', \xi_2{}', \xi_3{}'; x_1{}', x_2{}', x_3{}') = X_1. \quad \xi_1{}' = \xi_1, \xi_2{}' = \xi_3{}' \in \boldsymbol{R}, x_3{}' = x_3 \in \mathfrak{C}.$$

If $x_1{}' \neq 0$, define $\alpha_1(\pi x_1{}'/4|x_1{}'|) \in Spin_1(9)$. Then we get

$$\alpha_1(\pi x_1{}'/4|x_1{}'|)X_1 = (\xi_1{}'', \xi_2{}'', \xi_3{}''; 0, x_2{}'', x_3{}'') = X_2, \quad \xi_1{}'' = \xi_1{}', \xi_k{}'' \in \boldsymbol{R}, x_k{}'' \in \mathfrak{C}.$$

From the condition $X_2 \times X_2 = 0$, we have

$$\begin{array}{lll} \xi_2{}''\xi_3{}'' = 0, & \xi_3{}''\xi_1{}'' = x_2{}''\overline{x_2{}''}, & \xi_1{}''\xi_2{}'' = x_3{}''\overline{x_3{}''}, \\ \overline{x_2{}''}\,\overline{x_3{}''} = 0, & \xi_2{}''x_2{}'' = 0, & \xi_3{}''x_3{}'' = 0. \end{array}$$

Here we distinguish the following cases :

(I) When $\xi_2{}'' = 0$. From $X_2 \times X_2 = 0$, we have $x_3{}''\overline{x_3{}''} = 0$, hence $x_3{}'' = 0$. Therefore X_2 is of the form

$$X_2 = (\xi_1{}'', 0, \xi_3{}''; 0, x_2{}'', 0), \quad \xi_1{}'' = \xi_1{}', \xi_3{}'' \in \boldsymbol{R}, x_2{}'' \in \mathfrak{C}.$$

Choose $b \in \mathfrak{C}$ such that $(b, x_2{}'') = 0, |b| = \pi/4$, and define $\alpha_2(b) \in Spin_2(9)$ of Lemma 2.(2). Then

$$\alpha_2(b)X_2 = (\xi_1^{(3)}, 0, \xi_3^{(3)}; 0, x_2^{(3)}, 0) = X_3, \quad \xi_1^{(3)} = \xi_3^{(3)} \in \boldsymbol{R}, x_2^{(3)} \in \mathfrak{C}.$$

If $x_2^{(3)} \neq 0$, define $\alpha_2(\pi x_2^{(3)}/4|x_2^{(3)}|) \in Spin_2(9)$. Then

$$\alpha_2(\pi x_2^{(3)}/4|x_2^{(3)}|)X_3 = (\xi_1^{(4)}, 0, \xi_3^{(4)}; 0, 0, 0) = X_4, \quad \xi_1^{(4)}, \xi_3^{(4)} \in \boldsymbol{R}.$$

From $X_4 \times X_4 = 0$, we have $\xi_1^{(4)}\xi_3^{(4)} = 0$. If $\xi_1^{(4)} = 0$, define $\alpha_2(\pi/2) \in Spin_2(9)$. Then

$$\alpha_2(\pi/2)X_4 = (\xi_1^{(5)}, 0, 0; 0, 0, 0) = X_5, \quad \xi_1^{(5)} = \xi_3^{(4)} \in \boldsymbol{R}.$$

If $\xi_3^{(4)} = 0, X_4$ is also of the form X_5. Thus we obtain $X_5 = E_1$, since $\xi_1^{(5)} = \mathrm{tr}(X_5) = \mathrm{tr}(E_1) = 1$.

(II) When ${\xi_2}'' \neq 0$. From $X_2 \times X_2 = 0$, we have ${\xi_2}''{x_2}'' = 0$, hence ${x_2}'' = 0$. Therefore X_2 is of the form

$$X_2 = ({\xi_1}'', {\xi_2}'', 0; 0, 0, {x_3}''), \quad {\xi_k}'' \in \boldsymbol{R}, {x_3}'' \in \mathfrak{C}.$$

By considering $\alpha_1(\pi/2)X_2$, this can be reduced to the case (I).

We have just completed the proof of Theorem 5.

5 $Spin(10)$-generators of the group E_6

The simply connected compact Lie group E_6 is given by

$$E_6 = \{\alpha \in \mathrm{Iso}_C(\mathfrak{J}^C) \,|\, \alpha X \times \alpha Y = \tau\alpha\tau(X \times Y), < \alpha X, \alpha Y > = < X, Y >\}.$$

The group E_6 has subgroups

$$Spin_k(10) = \{\alpha \in E_6 \,|\, \alpha E_k = E_k\} \quad (k = 1, 2),$$

which is isomorphic to the usual spinor group $Spin(10)$ ([2],[3]).

Lemma 6. (1) *For any element*

$$X = (\xi_1, \xi_2, \xi_3; x_1, 0, 0), \quad \xi_k \in C, x_1 \in \mathfrak{C}^C$$

of $\mathfrak{J}^C$*, there exists some element* $\alpha_1 \in Spin_1(10)$ *such that*

$$\alpha_1 X = ({\xi_1}', {\xi_2}', {\xi_3}'; 0, 0, 0), \quad {\xi_1}' = \xi_1, {\xi_k}' \in C.$$

(2) *For any element*

$$X = (\xi_1, 0, 0; 0, x_2, x_3), \quad \xi_1 \in C, x_k \in \mathfrak{C}^C$$

of $\mathfrak{J}^C$*, there exists some element* $\alpha_1 \in Spin_1(9)$ *such that*

$$\alpha_1 X = (\xi_1, 0, 0; 0, {x_2}', {x_3}'), \quad {\xi_1}' = \xi_1 \in C, {x_2}' \in \mathfrak{C}^C, {x_3}' \in \mathfrak{C}.$$

Proof (1) For $x_1 = p + iq\,(p, q \in \mathfrak{C})$, choose $a \in \mathfrak{C}, a \neq 0$, such that $(a, p) = (a, q) = 0$, and define $\alpha_1(\pi a/4|a|) \in Spin_1(9)$ of Lemma 2.(1). Then

$$\alpha_1(\pi a/4|a|)X = ({\xi_1}', {\xi_2}', {\xi_2}' = {\xi_3}', {\xi_3}'; {x_1}', 0, 0) = X_1. \quad {\xi_1}' = \xi_1, {\xi_2}' = {\xi_3}' \in C, {x_1}' \in \mathfrak{C}^C.$$

Next, for ${x_1}' = p' + iq'\,(p', q' \in \mathfrak{C})$, choose $b \in \mathfrak{C}, b \neq 0$, such that $(b, p') = (b, q') = 0$, and define $\beta_1(\pi b/4|b|) \in Spin_1(10)$ of Lemma 3.(1). Then

$$\beta_1(\pi b/4|b|)X_1 = ({\xi_1}'', 0, 0; {x_1}'', 0, 0) = X_2, \quad {\xi_1}'' = \xi_1 \in C, {x_1}'' \in \mathfrak{C}^C.$$

Next, for ${x_1}'' = p'' + iq''$ $(p'', q'' \in \mathfrak{C})$, if $q'' \neq 0$, define $\alpha_1(\pi q''/4|q''|) \in Spin_1(9)$. Then

$$\alpha_1(\pi q''/4|q''|)X_2 = (\xi_1^{(3)}, \xi_2^{(3)}, \xi_3^{(3)}; p^{(3)}, 0, 0) = X_3, \quad \xi_1^{(3)} = \xi_1, \xi_3^{(3)} = -\xi_2^{(3)} \in C, p^{(3)} \in \mathfrak{C}.$$

Finally, if $p^{(3)} \neq 0$, define $\beta_1(\pi p^{(3)}/4|p^{(3)}|) \in Spin_1(10)$. Then we get

$$\beta_1(\pi p^{(3)}/4|p^{(3)}|)X_3 = (\xi_1^{(4)}, \xi_2^{(4)}, \xi_3^{(4)}; 0, 0, 0), \quad \xi_1^{(4)} = \xi_1, \xi_k^{(4)} \in C$$

as desired.

(2) At first, we show that for any element

$$Z = (\zeta_1, 0, 0; 0, z_2, z_3), \quad \zeta_1 \in \boldsymbol{R}, z_k \in \mathfrak{C},$$

there exists $\alpha_1 \in Spin_1(9)$ such that

$$\alpha_1 Z = (\zeta_1', 0, 0; 0, z_2', 0), \quad \zeta_1' \in \boldsymbol{R}, z_2' \in \mathfrak{C}.$$

In fact,if $z_2 z_3 \neq 0$, choose $t > 0$ such that $\cot(t|z_2 z_3|) = -|z_2|/|z_3|$, and define $\alpha_1(t\overline{z_2 z_3}) \in Spin_1(9)$. Then we get ($z_3$-part of $\alpha_1(t\overline{z_2 z_3})Z) = 0$. If $z_2 = 0$, then $\alpha_1(\pi/2)Z$ is of the form as desired.Now for a given element $X = (\xi_1, 0, 0; 0, x_2, x_3) \in \mathfrak{J}^C$, express $X = Y + iZ, Y, Z \in \mathfrak{J}$ and apply the result above to Z, then we get the required form $\alpha_1 X = \alpha_1 Y + i\alpha_1 Z$.

Theorem 7. *Any element $\alpha \in E_6$ can be represented as*

$$\alpha = \alpha_1 \alpha_2 {\alpha_1}', \quad \alpha_1, {\alpha_1}' \in Spin_1(10), \alpha_2 \in Spin_2(10).$$

Proof For a given element $\alpha \in E_6$, set $\alpha E_1 = (\xi_1, \xi_2, \xi_3; x_1, x_2, x_3) = X_0 \in \mathfrak{J}^C$. By Lemma 6.(1), we can take $\alpha_1 \in Spin_1(10)$ such that

$$\alpha_1 X_0 = ({\xi_1}', {\xi_2}', {\xi_3}'; 0, {x_2}', {x_3}') = X_1, \quad {\xi_1}' = \xi_1,$$

noting that the subspaces $\{(\xi_1, \xi_2, \xi_3; x_1, 0, 0) \in \mathfrak{J}^C\}$ and $\{(0, 0, 0; 0, x_2, x_3) \in \mathfrak{J}^C\}$ are invariant under the action of the elements of $Spin_1(10)$, respectively. From the condition $X_1 \times X_1 = 0$, we have ${\xi_2}'{\xi_3}' = 0$. As a result, the argumant is divided into the following three cases :

(I) Case ${\xi_2}' = 0, {\xi_3}' \neq 0$. From $X_1 \times X_1 = 0$, we have ${\xi_3}'{x_3}' = 0$, hence ${x_3}' = 0$. Therefore X_1 is of the form

$$X_1 = ({\xi_1}', 0, {\xi_3}'; 0, {x_2}', 0), \quad {\xi_1}' = \xi_1,$$

Thus, for $X_1 \in \mathfrak{J}^C$, we can take $\alpha_2 \in Spin_2(10)$ such that

$$\alpha_2 X_1 = ({\xi_1}'', 0, {\xi_3}''; 0, 0, 0) = X_2,$$

in the same way as in Lemma 6.(1). Then, from $X_2 \times X_2 = 0$, we have ${\xi_1}''{\xi_3}'' = 0$, tha is,

$$X_2 = ({\xi_1}'', 0, 0; 0, 0, 0), \ (\tau {\xi_1}''){\xi_1}'' = 1 \quad \text{or} \quad X_2 = (0, 0, {\xi_3}''; 0, 0, 0), \ (\tau {\xi_3}''){\xi_3}'' = 1.$$

Thus we obtain that there exist some elements $\varepsilon_2(t) \in Spin_2(10)$ and $\alpha_2(\pi/2) \in Spin_2(9)$ such that

$$\varepsilon_2(t)X_2 = E_1 \quad \text{or} \quad \varepsilon_2(t)\alpha_2(\pi/2)X_2 = E_1,$$

where $\varepsilon_2(t) \in Spin_2(10)$ is defined by

$$\varepsilon_2(t)(\xi_1, \xi_2, \xi_3; x_1, x_2, x_3) = (e^{it}\xi_1, \xi_2, e^{-it}\xi_3; e^{-it/2}x_1, x_2, e^{it/2}x_3), \quad t \in \boldsymbol{R}$$

(cf. [2] Lemma 10.(1)).

(II) Case ${\xi_2}' \neq 0, {\xi_3}' = 0$. From $X_1 \times X_1 = 0$, we have ${\xi_2}'{x_2}' = 0$, hence ${x_2}' = 0$. Therefore X_1 is of the form

$$X_1 = ({\xi_1}', {\xi_2}', 0; 0, 0, {x_3}'), \quad {\xi_1}' = \xi_1.$$

Thus, by considering $\alpha_1(\pi/2)X_1$, where $\alpha_1(\pi/2) \in Spin_1(9)$, this can be reduced to Case (I).

(III) Case ${\xi_2}' = {\xi_3}' = 0$. By Lemma 6.(2),we can take ${\alpha_1}' \in Spin_1(9)$ such that

$${\alpha_1}'X_1 = ({\xi_1}'', 0, 0; 0, {x_2}'', {x_3}'') = X_2, \quad {\xi_1}'' = \xi_1, {x_2}'' \in \mathfrak{C}^C, {x_3}'' \in \mathfrak{C}.$$

Then, from $X_2 \times X_2 = 0$ we have ${x_3}''\overline{x_3''} = 0$, hence ${x_3}'' = 0$. Thus, for $X_2 = ({\xi_1}'', 0, 0; 0, {x_2}'', 0) \in \mathfrak{J}^C$, we can take $\alpha_2 \in Spin_2(10)$ such that

$$\alpha_2 X_2 = (\xi_1^{(3)}, 0, \xi_3^{(3)}; 0, 0, 0) = X_3,$$

in the same way as in Lemma 6.(1). Hence in a similar way as in Case (I), we obtain the required result.

We have just completed the proof of Theorem 7.

6 $Spin(12)-$generators of the group E_7

The simply connected compact Lie group E_7 is given by

$$E_7 = \{\alpha \in \mathrm{Iso}_C(\mathfrak{P}^C) \,|\, \alpha(P \times Q)\alpha^{-1} = \alpha P \times \alpha Q, < \alpha P, \alpha Q > = < P, Q >\}.$$

The group E_7 has subgroups

$$Spin_k(12) = \{\alpha \in E_7 \,|\, \alpha\kappa_k = \kappa_k\alpha, \alpha\mu_k = \mu_k\alpha\} \quad (k = 1, 2)$$

where κ_k and μ_k are defined by

$$\begin{aligned} \kappa_k(X, Y, \xi, \eta) &= (-(E_k, X)E_k + 4E_k \times (E_k \times X), (E_k, Y)E_k - 4E_k \times (E_k \times Y), -\xi, \eta), \\ \mu_k(X, Y, \xi, \eta) &= (2E_k \times Y + \eta E_k, 2E_k \times X + \xi E_k, (E_k, Y), (E_k, X)), \end{aligned}$$

respectively, *e.g.*, when $k = 1$, for $P = ((\xi_1, \xi_2, \xi_3; x_1, x_2, x_3), (\eta_1, \eta_2, \eta_3; y_1, y_2, y_3), \xi, \eta) \in \mathfrak{P}^C$,

$$\begin{aligned}
&\kappa_1 P = ((-\xi_1, \xi_2, \xi_3; x_1, 0, 0), (\eta_1, -\eta_2, -\eta_3; -y_1, 0, 0), -\xi, \eta),\\
&\mu_1 P = ((\eta, \eta_3, \eta_2; -y_1, 0, 0), (\xi, \xi_3, \xi_2; -x_1, 0, 0), \eta_1, \xi_1).
\end{aligned}$$

Then $Spin_k(12)$ is isomorphic to the usual spinor group $Spin(12)$ ([2], [4]).

Lemma 8. *For an element* $P = ((\xi_1, \xi_2, \xi_3; x_1, x_2, x_3), (\eta_1, \eta_2, \eta_3; y_1, y_2, y_3), \xi, \eta) \in \mathfrak{P}^C$ *satisfying* $P \times P = 0$, *it holds the following*

$$(1) \quad \xi_1\eta_1 + \xi_2\eta_2 + \xi_3\eta_3 + 2(x_1, y_1) + 2(x_2, y_2) + 2(x_3, y_3) - 3\xi\eta = 0,$$

$$\begin{array}{llll}
(2) & \xi_2\xi_3 - \eta_1\eta - x_1\overline{x_1} = 0, & (3) & \xi_3\xi_1 - \eta_2\eta - x_2\overline{x_2} = 0,\\
(4) & \xi_1\xi_2 - \eta_3\eta - x_3\overline{x_3} = 0, & (5) & \xi_1 x_1 + \eta y_1 - \overline{x_2 x_3} = 0,\\
(6) & \xi_2 x_2 + \eta y_2 - \overline{x_3 x_1} = 0, & (7) & \xi_3 x_3 + \eta y_3 - \overline{x_1 x_2} = 0,\\
(8) & \eta_2\eta_3 - \xi_1\xi - y_1\overline{y_1} = 0, & (9) & \eta_3\eta_1 - \xi_2\xi - y_2\overline{y_2} = 0,\\
(10) & \eta_1\eta_2 - \xi_3\xi - y_3\overline{y_3} = 0, & (11) & \eta_1 y_1 + \xi x_1 - \overline{y_2 y_3} = 0,\\
(12) & \eta_2 y_2 + \xi x_2 - \overline{y_3 y_1} = 0, & (13) & \eta_3 y_3 + \xi x_3 - \overline{y_1 y_2} = 0,\\
(14) & \eta_3 x_1 + \xi_2 y_1 + \overline{y_2 x_3} = 0, & (15) & \eta_3 x_2 + \xi_1 y_2 + \overline{x_3 y_1} = 0,\\
(16) & \eta_2 x_3 + \xi_1 y_3 + \overline{y_1 x_2} = 0, & (17) & \eta_1 x_3 + \xi_2 y_3 + \overline{x_1 y_2} = 0.
\end{array}$$

Proof These are immediate from the straightforward computation of $P \times P = 0$. (Note that those are not all of the relations followed by $P \times P = 0$.)

Lemma 9. (1) *For any element* $P \in \mathfrak{P}^C$, *there exists some element* $\alpha_1 \in Spin_1(12)$ *such that*

$$\alpha_1 P = ((\xi_1, 0, 0; 0, x_2, x_3), (\eta_1, \eta_2, \eta_3; 0, y_2, y_3), \xi, \eta).$$

In particular, if an element $P = ((0, \xi_2, \xi_3; x_1, 0, 0), (\eta_1, 0, 0; 0, 0, 0), 0, \eta) \in \mathfrak{P}^C$ *satisfies the conditions* $P \times P = 0$ *and* $< P, P >= 1$, *then there exists some element* $\alpha_1 \in Spin_1(12)$ *such that*

$$\alpha_1 P = \underset{\cdot}{1}, \quad \textit{where } \underset{\cdot}{1} = (0, 0, 0, 1) \in \mathfrak{P}^C.$$

(2) *For any element* $P \in \mathfrak{P}^C$, *there exists some element* $\alpha_2 \in Spin_2(12)$ *such that*

$$\alpha_2 P = ((0, \xi_2, 0; x_1, 0, x_3), (\eta_1, \eta_2, \eta_3; y_1, 0, y_3), \xi, \eta).$$

In particular, if an element $P = ((\xi_1, 0, \xi_3; 0, x_2, 0), (0, \eta_2, 0; 0, 0, 0), 0, \eta) \in \mathfrak{P}^C$ *satisfies the conditions* $P \times P = 0$ *and* $< P, P >= 1$, *then there exists some element* $\alpha_2 \in Spin_2(12)$ *such that*

$$\alpha_2 P = \underset{\cdot}{1}.$$

Proof (1) The first half is the very [2] Proposition 4.(2). We shall now prove the latter half. For an element $P = ((0, \xi_2, \xi_3; x_1, 0, 0), (\eta_1, 0, 0; 0, 0, 0), 0, \eta) \in \mathfrak{P}^C$, act $\alpha_1 \in Spin_1(12)$ that is given in the first half which is composed of the elements of $Spin_1(12)$ defined in Lemmas 2, 3 and 4, on P. Then we get

$$\alpha_1 P = ((0, 0, 0; 0, 0, 0), ({\eta_1}', 0, 0; 0, 0, 0), 0, \eta') = P_1,$$

noting that the subspaces $< \mathfrak{P}^C >_1, < \mathfrak{P}^C >_1{}'$ and $< \mathfrak{P}^C >_1{}''$ of $\mathfrak{P}^C$ are invariant under the action of the elements of $Spin_1(12)$ defined in Lemmas 2, 3 and 4, respectively, where

$$< \mathfrak{P}^C >_1 = \{((\xi_1, 0, 0; 0, 0, 0), (0, \eta_2, \eta_3; y_1, 0, 0), \xi, 0) \in \mathfrak{P}^C\},$$

$$< \mathfrak{P}^C >_1{}' = \{((0, \xi_2, \xi_3; x_1, 0, 0), (\eta_1, 0, 0; 0, 0, 0), 0, \eta) \in \mathfrak{P}^C\},$$

$$< \mathfrak{P}^C >_1{}'' = \{((0, 0, 0; 0, x_2, x_3), (0, 0, 0; 0, y_2, y_3), 0, 0) \in \mathfrak{P}^C\}.$$

From $P \times P = 0$, we have $\eta_1{}'\eta' = 0$ by Lemma 8.(2). As a result, the argument is devided into the following three cases:

(I) Case $\eta_1{}' = 0, \eta' \neq 0. P_1$ is of the form $P_1 = ((0, 0, 0; 0, 0, 0), (0, 0, 0; 0, 0, 0), 0, \eta')$. Now, for $\theta \in C$ satifying $(\tau\theta)\theta = 1$, define the mapping $\epsilon_1(\theta) : \mathfrak{P}^C \to \mathfrak{P}^C$ as follows.

$$\begin{aligned}&\epsilon_1(\theta)((\xi_1, \xi_2, \xi_3; x_1, x_2, x_3), (\eta_1, \eta_2, \eta_3; y_1, y_2, y_3), \xi, \eta)\\ &\quad = ((\theta^{-2}\xi_1, \xi_2, \xi_3; x_1, \theta^{-1}x_2, \theta^{-1}x_3), (\theta^2\eta_1, \eta_2, \eta_3; y_1, \theta y_2, \theta y_3), \theta^2\xi, \theta^{-2}\eta).\end{aligned}$$

Then $\epsilon_1(\theta) \in Spin_1(12)$. Therefore, noting that $(\tau\eta')\eta' =< P_1, P_1 >= 1$, choose $\theta \in C$ such that $\theta^2 = \eta'$ and set $\epsilon_1(\theta)$. Then we get $\epsilon_1(\theta)P_1 = \underset{\cdot}{1}$.

(II) Case $\eta_1{}' \neq 0, \eta' = 0$. By considering $\gamma_1(\pi/2)P_1$, where $\gamma_1(\pi/2) \in Spin_1(12)$ of Lemma 4.(1), this can be reduced to Case (I).

(III) Case $\eta_1{}' = \eta' = 0$. This does not occur, because $< P_1, P_1 >= 1$.

(2) It is similarly verified by using $Spin_2(12)$ instead of $Spin_1(12)$ in the proof of (1).

Theorem 10. *Any element $\alpha \in E_7$ can be represented as*

$$\alpha = \alpha_1\alpha_2\alpha_1{}'\alpha_2{}'\alpha_1{}'', \quad \alpha_1, \alpha_1{}', \alpha_1{}'' \in Spin_1(12), \alpha_2, \alpha_2{}' \in Spin_2(12).$$

Proof For a given element $\alpha \in E_7$, it suffices to show that there exist $\alpha_1, \alpha_1{}' \in Spin_1(12)$ and $\alpha_2 \in Spin_2(12)$ such that $\alpha_1{}'\alpha_2\alpha_1\alpha\underset{\cdot}{1} = \underset{\cdot}{1}$. In fact, since an element $\alpha \in E_7$ belongs to $E_6 (\subset E_7)$ if and only if α fixes an alement $\underset{\cdot}{1}$, *i.e.,* $\alpha\underset{\cdot}{1} = \underset{\cdot}{1}$ ([4]), it follows $\alpha_1{}'\alpha_2\alpha_1\alpha \in E_6$, which implies that $\alpha \in E_7$ can be represented as a required form by Theorem 7. Now, set

$$\alpha\underset{\cdot}{1} = ((\xi_1, \xi_2, \xi_3; x_1, x_2, x_3), (\eta_1, \eta_2, \eta_3; y_1, y_2, y_3), \xi, \eta) = P_0 \in \mathfrak{P}^C.$$

Then, by Lemma 9.(1), we can take $\alpha_1 \in Spin_1(12)$ such that

$$\alpha_1 P_0 = ((\xi_1{}', 0, 0; 0, x_2{}', x_3{}'), (\eta_1{}', \eta_2{}', \eta_3{}'; 0, y_2{}', y_3{}'), \xi', \eta') = P_1.$$

From $P_1 \times P_1 = 0$, we have $\eta_1{}'\eta' = 0$ by Lemma 8.(2). As a result, the argument is devided into the following three cases :

(I) Case $\eta_1{}' = 0, \eta' \neq 0$. By Lemma 8.(6) and (7), we get $y_2{}' = y_3{}' = 0$. Furthermore we get $\xi' = 0$ by Lemma 8.(1). Therefore P_1 is of the form

$$P_1 = ((\xi_1{}', 0, 0; 0, x_2{}', x_3{}'), (0, \eta_2{}', \eta_3{}'; 0, 0, 0), 0, \eta').$$

Then, by Lemma 8.(8), we have ${\eta_2}'{\eta_3}' = 0$. Hence there are three cases to be considered.

(I.A) Case ${\eta_2}' = 0, {\eta_3}' \neq 0$. By Lemma 8.(15), we get ${x_2}' = 0$, that is, P_1 is of the form

$$P_1 = (({\xi_1}', 0, 0; 0, 0, {x_3}'), (0, 0, {\eta_3}'; 0, 0, 0), 0, \eta').$$

Then, applying Lemma 9.(2) to $\alpha_1(\pi/2)P_1$, where $\alpha_1(\pi/2) \in Spin_1(9)$, we can obtain that there exists some element $\alpha_2 \in Spin_2(12)$ such that $\alpha_2\alpha_1(\pi/2)P_1 = \underset{\cdot}{1}$.

(I.B) Case ${\eta_2}' \neq 0, {\eta_3}' = 0$. By Lemma 8.(16), we get ${x_3}' = 0$, that is, P_1 is of the form

$$P_1 = (({\xi_1}', 0, 0; 0, {x_2}', 0), (0, {\eta_2}', 0; 0, 0, 0), 0, \eta').$$

Thus we can easily obtain the required result by Lemma 9.(2).

(I.C) Case ${\eta_2}' = {\eta_3}' = 0$. P_1 is of the form

$$P_1 = (({\xi_1}', 0, 0; 0, {x_2}', {x_3}'), (0, 0, 0; 0, 0, 0), 0, \eta').$$

Here we distinguish the following cases :

(I.C.1) When ${x_2}' \neq 0, {x_3}' \neq 0$. By Lemma 6.(2), we can take ${\alpha_1}' \in Spin_1(9)$ such that

$${\alpha_1}'P_1 = (({\xi_1}'', 0, 0; 0, {x_2}'', {x_3}''), (0, 0, 0; 0, 0, 0), 0, \eta''), \quad {\xi_1}'' = {\xi_1}', \eta'' = \eta' \in C, {x_2}'' \in \mathfrak{C}^C, {x_3}'' \in \mathfrak{C}.$$

Then, by Lemma 8.(4) we have ${x_3}''\overline{{x_3}''} = 0$, hence ${x_3}'' = 0$. Thus we easily obtain the required result by Lemma 9.(2).

(I.C.2) When ${x_2}' = 0, {x_3}' \neq 0$. Considering $\alpha_1(\pi/2)P_1$, where $\alpha_1(\pi/2) \in Spin_1(9)$, we can easily obtain the required result by Lemma 9.(2).

(I.C.3) When ${x_2}' \neq 0, {x_3}' = 0$. We can easily obtain the required result by Lemma 9.(2).

(I.C.4) When ${x_2}' = {x_3}' = 0$. We can easily obtain the required result by Lemma 9.(2).

(II) Case ${\eta_1}' \neq 0, \eta' = 0$. By considering $\delta_1(\pi/2)P_1$, where $\delta_1(\pi/2) \in Spin_1(12)$ of Lemma 4.(3), this can be reduced to Case (I).

(III) Case ${\eta_1}' = \eta' = 0$. P_1 is of the form

$$P_1 = (({\xi_1}', 0, 0; 0, {x_2}', {x_3}'), (0, {\eta_2}', {\eta_3}'; 0, {y_2}', {y_3}'), \xi', 0).$$

Now, as is similar to Lemma 9.(1), we obtain that, for any element $P \in \mathfrak{P}^C$, there exists some element $\alpha_1 \in Spin_1(12)$ such that

$$\alpha_1 P = ((\xi_1, \xi_2, \xi_3; 0, x_2, x_3), (\eta_1, 0, 0; 0, y_2, y_3), \xi, \eta).$$

Note that the invariant subspaces $< \mathfrak{P}^C >_1, < \mathfrak{P}^C >_1{}'$ and $< \mathfrak{P}^C >_1{}''$ of $\mathfrak{P}^C$ under the action of the elements of $Spin_1(12)$ defined in Lemmas 2, 3 and 4.

Then, applying the result above to the present case (III), we can take ${\alpha_1}' \in Spin_1(12)$ such that

$${\alpha_1}'P_1 = (({\xi_1}'', 0, 0; 0, {x_2}'', {x_3}''), (0, 0, 0; 0, {y_2}'', {y_3}''), \xi'', 0) = P_2.$$

Therefore we have ${\xi_1}''\xi'' = 0$ by Lemma 8.(8). Hence there are three cases to be considered.

(III.A) Case ${\xi_1}'' = 0, \xi'' \neq 0$. By Lemma 8.(12) and (13), we get ${x_2}'' = {x_3}'' = 0$. Then P_2 is of the form

$$P_2 = ((0, 0, 0; 0, 0, 0), (0, 0, 0; 0, {y_2}'', {y_3}''), \xi'', 0).$$

Thus, by considering $\gamma_1(\pi/2)P_2$, where $\gamma_1(\pi/2) \in Spin_1(12)$ of Lemma 4.(1), this can be reduced to Case (I.C).

(III.B) Case ${\xi_1}'' \neq 0, \xi'' = 0$. By Lemma 8.(15) and (16), we get ${y_2}'' = {y_3}'' = 0$. Therefore this is reduced to Case (I.C).

(III.C) Case ${\xi_1}'' = \xi'' = 0$. P_2 is of the form

$$P_2 = ((0, 0, 0; 0, {x_2}'', {x_3}''), (0, 0, 0; 0, {y_2}'', {y_3}''), 0, 0).$$

Here we distinguish the following cases :

(III.C.1) When ${x_2}'' \neq 0$. By Lemma 9.(2), there exists some element $\alpha_2 \in Spin_2(12)$ such that

$$\alpha_2 P_2 = ((0, \xi_2^{(3)}, 0; x_1^{(3)}, 0, x_3^{(3)}), (\eta_1^{(3)}, \eta_2^{(3)}, \eta_3^{(3)}; y_1^{(3)}, 0, y_3^{(3)}), \xi^{(3)}, \eta^{(3)}) = P_3.$$

Here, by Lemma 8.(3), we have $\eta_2^{(3)}\eta^{(3)} = 0$. Hence there are three cases to be considered.

(III.C.1.1) Case $\eta_2^{(3)} = 0, \eta^{(3)} \neq 0$. By Lemma 8.(5) and (7), we get $y_1^{(3)} = y_3^{(3)} = 0$. Furthermore, we get $\xi^{(3)} = 0$ by Lemma 8.(1). Then P_3 is of the form

$$P_3 = ((0, \xi_2^{(3)}, 0; x_1^{(3)}, 0, x_3^{(3)}), (\eta_1^{(3)}, 0, \eta_3^{(3)}; 0, 0, 0), 0, \eta^{(3)}).$$

Here, by lemma 8.(9), we have $\eta_3^{(3)}\eta_1^{(3)} = 0$. Hence there are three cases to be considered.

(III.C.1.1.1) Case $\eta_1^{(3)} = 0, \eta_3^{(3)} \neq 0$. By Lemma 8.(14), we get $x_1^{(3)} = 0$. Then, considering $\alpha_2(\pi/2)P_3$, where $\alpha_2(\pi/2) \in Spin_2(9)$, we can easily obtain the required result by Lemma 9.(1).

(III.C.1.1.2) Case $\eta_1^{(3)} \neq 0, \eta_3^{(3)} = 0$. By Lemma 8.(17), we get $x_3^{(3)} = 0$. Then we can easily obtain the required result by Lemma 9.(1).

(III.C.1.1.3) Case $\eta_1^{(3)} = \eta_3^{(3)} = 0$. P_3 is of the form

$$P_3 = ((0, \xi_2^{(3)}, 0; x_1^{(3)}, 0, x_3^{(3)}), (0, 0, 0; 0, 0, 0), 0, \eta^{(3)}).$$

Here we distinguish the following cases :

(III.C.1.1.3.(i)) When $x_1^{(3)} \neq 0, x_3^{(3)} \neq 0$. As is similar to Lemma 6.(2), we obtain that there exists some element ${\alpha_2}' \in Spin_2(9)$ such that

$$\begin{aligned}\alpha_2{}'P_3 = ((0, \xi_2^{(4)}, 0; x_1^{(4)}, 0, x_3^{(4)}), (0,0,0;0,0,0), 0, \eta^{(4)}) = P_4,\\ \xi_2^{(4)} = \xi_2^{(3)}, \eta^{(4)} = \eta^{(3)} \in C,\\ x_1^{(4)} \in \mathfrak{C}, x_3^{(4)} \in \mathfrak{C}^C.\end{aligned}$$

Then, by Lemma 8.(2), we have $x_1^{(4)}\overline{x_1^{(4)}} = 0$, hence $x_1^{(4)} = 0$. Thus, considering $\alpha_2(\pi/2)P_4$, where $\alpha_2(\pi/2) \in Spin_2(9)$, we can easily obtain the required result by Lemma 9.(1).

(III.C.1.1.3.(ii)) When $x_1^{(3)} = 0, x_3^{(3)} \neq 0$. Considering $\alpha_2(\pi/2)P_3$, where $\alpha_2(\pi/2) \in Spin_2(9)$, we can easily obtain the required result by Lemma 9.(1).

(III.C.1.1.3.(iii)) When $x_1^{(3)} \neq 0, x_3^{(3)} = 0$. We easily obtain the required result by Lemma 9.(1).

(III.C.1.1.3.(iv)) When $x_1^{(3)} = x_3^{(3)} = 0$. We easily obtain the required result by Lemma 9.(1).

(III.C.1.2) Case $\eta_2^{(3)} \neq 0, \eta^{(3)} = 0$. By considering $\gamma_2(\pi/2)P_3$, where $\gamma_2(\pi/2) \in Spin_2(12)$ of Lemma 4.(2), this can be reduced to Case (III.C.1.1).

(III.C.1.3) Case $\eta_2^{(3)} = \eta^{(3)} = 0$. This does not occur. In fact, note that the subspace $< \mathfrak{P}^C >_2$ of $\mathfrak{P}^C$ is invariant under the action of the elements of $Spin_2(12)$ defined in Lemmas 2, 3 and 4, where $< \mathfrak{P}^C >_2 = \{((\xi_1, 0, \xi_2, ; 0, x_2, 0), (0, \eta_2, 0; 0, 0, 0), 0, \eta) \in \mathfrak{P}^C\}$. Then, for $P_3 = \alpha_2 P_2$, that is,

$$\begin{aligned}&((0, \xi_2^{(3)}, 0; x_1^{(3)}, 0, x_3^{(3)}), (\eta_1^{(3)}, 0, \eta_3^{(3)}; y_1^{(3)}, 0, y_3^{(3)}), \xi^{(3)}, 0)\\ &\quad = \alpha_2((0,0,0;0,x_2{}'', x_3{}''), (0,0,0;0,y_2{}'', y_3{}''), 0, 0),\end{aligned}$$

where $\alpha_2 \in Spin_2(12)$, the condition $\eta_2^{(3)} = \eta^{(3)} = 0$ contradicts $x_2{}'' \neq 0$.

(III.C.2) When $x_2{}'' = 0, x_3{}'' \neq 0$. By considering $\alpha_1(\pi/2)P_2$, where $\alpha_1(\pi/2) \in Spin_1(9)$, this can be reduced to Case (III.C.1).

(III.C.3) When $x_2{}'' = x_3{}'' = 0, y_3{}'' \neq 0$. By considering $\gamma_1(\pi/2)P_2$, where $\gamma_1(\pi/2) \in Spin_1(12)$, this can be reduced to Case (III.C.1).

(III.C.4) When $x_2{}'' = x_3{}'' = y_3{}'' = 0, y_2{}'' \neq 0$. By considering $\alpha_1(\pi/2)P_2$, where $\alpha_1(\pi/2) \in Spin_1(9)$, this can be reduced to Case (III.C.3).

(III.C.5) When $x_2{}'' = x_3{}'' = y_2{}'' = y_3{}'' = 0$. It is obvious that this does not occur.

We have just completed the proof of Theorem 10.

Conjecture We know that the simply connected compact excetional Lie group E_8 has subgroups $Ss_k(16) = (E_8)^{\sigma_k}$ (where $\sigma_k = \exp \pi\kappa_k$), $k = 1, 2, 3$ (which is isomorphic to $Spin(16)/\boldsymbol{Z}_2$ not $SO(16)$). Now the authors do not know if $Ss_1(16)$ and $Ss_2(16)$ generate the group E_8?

References

[1] C. Chevalley, *Theorey of Lie groups*, Princeton Univ. Press, New Jersey, 1946.

[2] T. Miyasaka, O. Yasukura and I. Yokota, *Diagonalization of an element* P *of* $\mathfrak{P}^C$ *by the compact Lie group* E_7, Tsukuba J.Math., 22(1998), 687-703.

[3] I. Yokota, *Realization of involutive automorphisms* σ *and* G^σ *of exceptional linear Lie groups* G, Part I, $G = G_2, F_4$ *and* E_6, Tsukuba J. Math., 14(1990), 185-223.

[4] I. Yokota, *Realization of involutive automorphisms* σ *and* G^σ *of exceptional linear Lie groups* G, Part II, $G = E_7$, Tsukuba J. Math., 14(1990), 379-404.